arXiv:1701.02616v1 [math.AP] 10 Jan 2017

INTEGRAL ESTIMATES OF CONFORMAL DERIVATIVES AND SPECTRAL PROPERTIES OF THE NEUMANN-LAPLACIAN

V. GOL'DSHTEIN, V. PCHELINTSEV, A. UKHLOV

Abstract. In this paper we study integral estimates of derivatives of conformal mappings $\varphi : \mathbb{D} \to \Omega$ of the unit disc $\mathbb{D} \subset \mathbb{C}$ onto bounded domains Ω that satisfy the Ahlfors condition.These integral estimates lead to estimates of constants in Sobolev-Poincaré inequalities and by the Rayleigh quotient we obtain spectral estimates of the Neumann-Laplace operator in non-Lipschitz domains (quasidiscs) in terms of the (quasi)conformal geometry of the domains. The lower estimates of the first non-trivial eigenvalues of the Neumann-Laplace operator in some fractal type domains (snowflakes) were obtained.

1. Introduction

1.1. **Estimates of Conformal Derivatives.** In the work [19] we obtained lower estimates of the first non-trivial eigenvalues of the Neumann-Laplace operator in the terms of integrals of complex derivatives (i. e. hyperbolic metrics) of conformal mappings $\varphi : \mathbb{D} \to \Omega$. Let us recall that the classical Koebe distortion theorem [7] gives the following estimates of the complex derivatives in the case of univalent analytic functions (conformal homeomorphisms): $\varphi : \mathbb{D} \to \Omega$ normalized so that $\varphi(0) = 0$ and $\varphi'(0) = 1$:

$$\frac{1-|z|}{(1+|z|)^3} \leq |\varphi'(z)| \leq \frac{1+|z|}{(1-|z|)^3}.$$

The example of the complex plane slit along the negative real axis shows that these estimates don't give even square integrability of the complex derivatives in arbitrary simply connected planar domains. But if $\Omega \subset \mathbb{C}$ is a simply connected planar domain of finite measure then by simple calculation

$$\iint\limits_{\mathbb{D}} |\varphi'(z)|^2 \, dxdy = \iint\limits_{\mathbb{D}} J(z,\varphi) \, dxdy = |\Omega| < \infty.$$

(We identify the complex plane $\mathbb{C}$ and the real plane $\mathbb{R}^2$: $\mathbb{C} \ni z = x + iy = (x, y) \in \mathbb{R}^2$.)

Hence in special classes of domains we have better integral estimates of the complex derivatives than by the Koebe distortion theorem.

In the present work we study integral estimates of the complex derivatives in domains bounded by Jordan curves that satisfy to the Ahlfors three points condition

[0]**Key words and phrases:** Sobolev spaces, conformal mappings, quasiconformal mappings, elliptic equations.

[0]**2010 Mathematics Subject Classification:** 35P15, 46E35, 30C65.

[3]. For this study we introduce a notion of hyperbolic (integral) α-dilatation of Ω:

$$Q(\alpha,\Omega):=\iint\limits_{\mathbb{D}}|\varphi'(z)|^{\alpha}\,dxdy=\iint\limits_{\Omega}\left|\left(\varphi^{-1}\right)'(w)\right|^{2-\alpha}dudv.$$

The hyperbolic α-dilatation and its convergence hyperbolic interval

$$\mathrm{HI}(\Omega):=\{\alpha\in\mathbb{R}:Q(\alpha,\Omega)<\infty\}.$$

do not depend on choice of a conformal mapping $\varphi:\mathbb{D}\to\Omega$ and can be reformulated in terms of the hyperbolic metrics [10]. Namely

$$\begin{aligned}\iint\limits_{\mathbb{D}}|\varphi'(z)|^{\alpha}\,dxdy&=\iint\limits_{\mathbb{D}}\left(\frac{\lambda_{\mathbb{D}}(z)}{\lambda_{\Omega}(\varphi(z))}\right)^{\alpha}dxdy\\&=\iint\limits_{\Omega}\left|\left(\varphi^{-1}\right)'(w)\right|^{2-\alpha}dudv=\iint\limits_{\Omega}\left(\frac{\lambda_{\mathbb{D}}\left(\varphi^{-1}(w)\right)}{\lambda_{\Omega}(w)}\right)^{2-\alpha}dudv\end{aligned}$$

where $\lambda_{\mathbb{D}}$ and λ_{Ω} are hyperbolic metrics in $\mathbb{D}$ and Ω [6]. Let us recall that $\lambda_{\mathbb{D}}\left(\varphi^{-1}(w)\right)$ does not depends on choice of a conformal homeomorphism $\varphi:\Omega\to\mathbb{D}$ because any other conformal homeomorphism $\psi^{-1}:\Omega\to\mathbb{D}$ is a composition of φ^{-1} and a Möbius homeomorphism (that is an isometry of the hyperbolic metric).

Remark 1.1. For any bounded simply connected domain $(-1.78,2]\subset\mathrm{HI}(\alpha,\Omega)$ [18, 20]. A more detailed discussion about the hyperbolic α-dilatation and its convergence hyperbolic interval can be found in Appendix.

In [18] we proved that if a number $\alpha>2$ belongs to $\mathrm{HI}(\Omega)$ then Ω has finite geodesic diameter. By this reason we call domains that satisfy to a property $2<\alpha\in\mathrm{HI}(\Omega)$ as conformal α-regular domains [10].

In this paper we obtain estimates of $Q(\alpha,\Omega)$ for a large class of conformal α-regular domains (so-called quasidiscs) with the help of the inverse Hölder inequality for Jacobians of quasiconformal mappings. Recall that quasidiscs are images of the unit discs under quasiconformal homeomorphisms of the complex plane. This class includes all Lipschitz simply connected domains but also includes a class of fractal domains (for example, so-called Rohde snowflakes). Hausdorff dimensions of quasidiscs boundary can be any number of $[1,2)$.

With the help of the estimates for $Q(\alpha,\Omega)$ we obtain estimates of constants for Sobolev-Poincaré inequalities and as a result we obtain lower estimates for first nontrivial eigenvalues of the Laplace operator with the Neumann boundary condition.

1.2. **Spectral Estimates of the Neumann-Laplacian.** The weak formulation of the spectral problem for the Laplace operator with the Neumann boundary condition is the following: a function u solves this spectral problem iff $u\in W_2^1(\Omega)$ and

$$\iint\limits_{\Omega}\nabla u(x,y)\cdot\nabla v(x,y)\,dxdy=\mu\iint\limits_{\Omega}u(x,y)v(x,y)\,dxdy$$

for all $v\in W_2^1(\Omega)$.

Let us give a short historical remark about eigenvalues estimates for the Neumann-Laplace operator. In 1961 G.Polya [25] obtained upper estimates for eigenvalues in

so-called plane-covering domains. Namely, for the first nontrivial eigenvalue $\mu_1(\Omega)$ it is:

$$\mu_1(\Omega) \le 4\pi|\Omega|^{-1}.$$

The lower estimates for the $\mu_1(\Omega)$ for convex domains were obtained in the classical work [24]. It was proved that if Ω is convex with diameter $d(\Omega)$ (see, also [11, 14, 28]), then

$$\mu_1(\Omega) \ge \frac{\pi^2}{d(\Omega)^2}.$$

In [19] was proved, that if $\Omega \subset \mathbb{C}$ be a conformal α-regular domain, then the spectrum of Neumann-Laplace operator in Ω is discrete, can be written in the form of a non-decreasing sequence

$$0 = \mu_0(\Omega) < \mu_1(\Omega) \le \mu_2(\Omega) \le ... \le \mu_n(\Omega) \le ...,$$

and

$$1/\mu_1(\Omega) \le \frac{4}{\sqrt[\alpha]{\pi^2}} \left(\frac{2\alpha-2}{\alpha-2}\right)^{\frac{2\alpha-2}{\alpha}} Q(\alpha,\Omega)^{2/\alpha}. \tag{1.1}$$

Specifying by means of the inverse Hölder inequality and the Ahlfors condition this estimate for quasidiscs we obtain discreetness of the spectrum of the Neumann-Laplace operator in so-called Rohde snowflake domains and gives the lower estimate of first non-trivial eigenvalues:

Theorem A. *Let $S_p \subset \mathbb{C}$, $1/4 \le p < 1/2$, be the Rohde snowflake. Then the spectrum of Neumann-Laplace operator in S_p is discrete, can be written in the form of a non-decreasing sequence*

$$0 = \mu_0(S_p) < \mu_1(S_p) \le \mu_2(S_p) \le ... \le \mu_n(S_p) \le ...,$$

and

$$\frac{1}{\mu_1(S_p)} \le \frac{C_\alpha^2 e^{4\left(1+e^{2\pi(16/(1-2p))^5}\right)}}{2^{40}\pi} \left(\frac{2\alpha-2}{\alpha-2}\right)^{\frac{2\alpha-2}{\alpha}} \times \exp\left\{\frac{\pi^2(2+\pi^2)^2 e^{4\left(1+e^{2\pi(16/(1-2p))^5}\right)}}{2^{41}\log 3}\right\} |S_p|,$$

for $2 < \alpha < \frac{2K^2}{K^2-1}$, where

$$C_\alpha = \frac{10^6}{[(\alpha-1)(1-\nu)]^{1/\alpha}}, \quad \nu = 10^{4\alpha}\left(\frac{3\pi^2}{2^{37}} e^{4\left(1+e^{2\pi C^5}\right)}\right)^\alpha \frac{\alpha-2}{\alpha-1} < 1.$$

The suggested approach is based on estimates of hyperbolic metric in quasidiscs. Recall that a domain $\Omega \subset \mathbb{C}$ is called a K-quasidisc if it is the image of the unit disc $\mathbb{D}$ under a K-quasiconformal homeomorphism of the complex plane $\mathbb{C}$ onto itself. Note that quasidiscs represent large class domains including fractal type domains like snowflakes.

Follow [4] a homeomorphism $\varphi : \Omega \to \Omega'$ between planar domains is called K-quasiconformal if it preserves orientation, belongs to the Sobolev class $W^1_{2,\mathrm{loc}}(\Omega)$ and its directional derivatives ∂_ξ satisfy the distortion inequality

$$\max_\xi |\partial_\xi \varphi| \le K \min_\xi |\partial_\xi \varphi| \text{ a.e. in } \Omega.$$

If Ω is a K-quasidisc, then a conformal mapping $\varphi : \mathbb{D} \to \Omega$ allows K^2-quasiconformal reflection. It is well known that Jacobians of quasiconformal mappings satisfy the weak inverse Hölder inequality [8]. On the base of the weak inverse Hölder inequality and the estimates of the constants in doubling conditions for measures generated by Jacobians of quasiconformal mappings (Proposition 3.6) we obtain

Theorem C. *Let $\Omega \subset \mathbb{C}$ be a K-quasidisc. Then the spectrum of Neumann-Laplace operator in Ω is discrete, can be written in the form of a non-decreasing sequence*

$$0 = \mu_0(\Omega) < \mu_1(\Omega) \leq \mu_2(\Omega) \leq ... \leq \mu_n(\Omega) \leq ...,$$

and

$$\frac{1}{\mu_1(\Omega)} \leq \frac{K^2 C_\alpha^2}{\pi} \left(\frac{2\alpha - 2}{\alpha - 2}\right)^{\frac{2\alpha-2}{\alpha}} \exp\left\{\frac{K^2\pi^2(2+\pi^2)^2}{2\log 3}\right\} \cdot |\Omega|, \tag{1.2}$$

for $2 < \alpha < \frac{2K^2}{K^2-1}$, where

$$C_\alpha = \frac{10^6}{[(\alpha-1)(1-\nu)]^{1/\alpha}}, \quad \nu = 10^{4\alpha}\frac{\alpha-2}{\alpha-1}\left(24\pi^2K^2\right)^\alpha < 1.$$

The main technical problem of this estimate is evaluations of the quasiconformality coefficient K for quasidiscs. For this aim we use an equivalent description of quasidiscs in the terms of the Ahlfors's 3-point condition. This description of quasidiscs allows to obtain the estimates for fractal type domains (Theorem A).

On the base Theorem A and Theorem C we can assert that eigenvalues of the Laplace operator depend on quasi(conformal) geometry of planar domains.

2. Sobolev spaces and Poincaré-Sobolev inequalities

Let $E \subset \mathbb{C}$ be a measurable set. For any $1 \leq p < \infty$ we consider the Lebesgue space of locally integrable functions with the finite norm

$$\|f \mid L_p(E)\| := \left(\iint\limits_E |f(x,y)|^p \, dxdy\right)^{1/p} < \infty.$$

We define the Sobolev space $W_p^1(\Omega)$, $1 \leq p < \infty$, as a Banach space of locally integrable weakly differentiable functions $f : \Omega \to \mathbb{R}$ equipped with the following norm:

$$\|f \mid W_p^1(\Omega)\| = \left(\iint\limits_\Omega |f(x,y)|^p \, dxdy\right)^{\frac{1}{p}} + \left(\iint\limits_\Omega |\nabla f(x,y)|^p \, dxdy\right)^{\frac{1}{p}}.$$

We also define the homogeneous seminormed Sobolev space $L_p^1(\Omega)$, $1 \leq p < \infty$, of locally integrable weakly differentiable functions $f : \Omega \to \mathbb{R}$ equipped with the following seminorm:

$$\|f \mid L_p^1(\Omega)\| = \left(\iint\limits_\Omega |\nabla f(x,y)|^p \, dxdy\right)^{\frac{1}{p}}.$$

Recall that the embedding operator $i : L_p^1(\Omega) \to L_{1,\mathrm{loc}}(\Omega)$ is bounded.

If $f \in W_p^1(\mathbb{D})$, $1 \le p < \infty$, then for $0 \le \kappa = 1/p - 1/q < 1/2$ the Poincaré-Sobolev inequality

$$\left(\iint\limits_{\mathbb{D}} |f(z) - f_{\mathbb{D}}|^q \, dxdy \right)^{\frac{1}{q}} \le B_{q,p}(\mathbb{D}) \left(\iint\limits_{\mathbb{D}} |\nabla f(z)|^p \, dxdy \right)^{\frac{1}{p}}$$

holds (see, for example, [15, 19]) with the constant

$$B_{q,p}(\mathbb{D}) \le \frac{2}{\pi^\kappa} \left(\frac{1-\kappa}{1/2-\kappa} \right)^{1-\kappa}.$$

This estimate does not applicable in the critical case $p = 1$ and $q = 2$. Now we obtain the upper estimate of the Poincaré constant in the Poincaré-Sobolev inequality for the Sobolev space $W_1^1(\mathbb{D})$ in this critical case. We use the following Gagliardo inequality for functions with compact support [13, 23]:

$$\left(\iint\limits_{\Omega} \left| f(z) \right|^2 \, dxdy \right)^{\frac{1}{2}} \le \frac{1}{2\sqrt{\pi}} \iint\limits_{\Omega} |\nabla f(z)| \, dxdy,$$

where $\Omega \subset \mathbb{C}$ be a bounded Lipschitz domain.

Theorem 2.1. *Let $f \in W_1^1(\Omega)$. Then for any $r > 0$ and any $z_0 \in \Omega : \operatorname{dist}(z_0, \partial\Omega) > 2r$, the following inequality*

$$\left(\iint\limits_{D(z_0,r)} |f(z) - f_{D(z_0,r)}|^2 \, dxdy \right)^{\frac{1}{2}} \le \frac{3\sqrt{\pi^3}}{4} \iint\limits_{D(z_0,r)} |\nabla f(z)| \, dxdy \tag{2.1}$$

holds.

Proof. We prove this inequality in the case $f_{D(z_0,r)} = 0$. In this case the inequity (2.1) can be rewritten as

$$\left(\iint\limits_{D(z_0,r)} |f(z)|^2 \, dxdy \right)^{\frac{1}{2}} \le \frac{3\sqrt{\pi^3}}{4} \iint\limits_{D(z_0,r)} |\nabla f(z)| \, dxdy. \tag{2.2}$$

The inequality (2.2) is invariant under translations and similarities and it is sufficient to prove one in for the disc $D(0,1)$.

Denote by $D(0,\delta)$ an open disc of the radius $\delta > 1$. Choose the cut function η in the form $\eta = \eta(|z|)$ such that

$$\eta(z) = \begin{cases} 1, & \text{if } |z| < 1, \\ 0, & \text{if } |z| > \delta \end{cases}$$

and linear for $z \in R_\delta$, where $R_\delta := D(0,\delta) \setminus \overline{D(0,1)}$. Then

$$|\nabla \eta(z)| = \begin{cases} \frac{\delta}{\delta - 1} & \text{if } z \in R_\delta, \\ 0 & \text{otherwise.} \end{cases}$$

Let $\widetilde{f} : R_\delta \to \mathbb{R}$ be the extension function is defined by the rule

$$\widetilde{f}(z) = f\left(w(z)\right) \eta(z), \ w \in R_\delta,$$

where $w(z) = 1/\overline{z}$ is the inversion about the unit circle $S(0,1)$.

Define the extension operator on Sobolev spaces

$$E : L_1^1(D(0,1)) \to L_1^1(D(0,\delta))$$

by the formula

$$(Ef)(z) = \begin{cases} f(z) & \text{if } z \in D(0,1), \\ \widetilde{f}(z) & \text{if } z \in R_\delta. \end{cases}$$

In order to estimate the norm $\|E\|$ of the extension operator $E : L_1^1(D(0,1)) \to L_1^1(D(0,\delta))$ we have

$$\|E \mid L_1^1(\mathbb{D}(0,\delta))\| = \iint\limits_{D(0,1)} |\nabla f(z)| \, dxdy + \iint\limits_{R_\delta} |\nabla \widetilde{f}(z)| \, dxdy.$$

For estimate of the second integral in the right side of this equality by elementary calculations we have

$$|\nabla \widetilde{f}(z)| = |\nabla \left(f(w(z))\eta(z)\right)| = |\nabla \left(f(w(z))\right) \cdot \eta(z) + f(w(z)) \cdot \nabla\eta(z)| \leq |\nabla f(w(z))| + \frac{\delta}{\delta - 1} |f(w(z))|. \tag{2.3}$$

Hence

$$\iint\limits_{R_\delta} |\nabla \widetilde{f}(w(z))| \, dxdz \leq \iint\limits_{R_\delta} |\nabla f(w(z))| \, dxdy + \frac{\delta}{\delta - 1} \iint\limits_{R_\delta} |f(w(z))| \, dxdy.$$

First consider the integral

$$\iint\limits_{R_\delta} |\nabla f(w(z))| \, dxdy = \iint\limits_{R_\delta} |\nabla f|(w(z))|w'(z)|dxdy =$$

$$= \iint\limits_{R_\delta} |\nabla f|(w(z))|w'(z)||J(z,w)||J(z,w)|^{-1}dxdy$$

$$\leq \sup_{z \in R_\delta} \frac{|w'(z)|}{|J(z,w)|} \iint\limits_{R_\delta} |\nabla f|(w(z))|J(z,w)| \, dxdy$$

$$= \sup_{z \in R_\delta} \frac{|w'(z)|}{|J(z,w)|} \iint\limits_{D_\delta} |\nabla f|(w) \, dudv,$$

where $w : R_\delta \to D_\delta$, $w(z) = 1/\overline{z}$ and $D_\delta = \{w \in \mathbb{C} : 1/\delta < |z| < 1\}$. We calculate the norm of the derivative of mapping w by the formula

$$|w'(z)| = |w_z| + |w_{\overline{z}}|$$

and the Jacobian of mapping w by the formula

$$J(z,w) = |w_z|^2 - |w_{\overline{z}}|^2.$$

Here

$$w_z = \frac{1}{2}\left(\frac{\partial w}{\partial x} - i\frac{\partial w}{\partial y}\right) \quad \text{and} \quad w_{\overline{z}} = \frac{1}{2}\left(\frac{\partial w}{\partial x} + i\frac{\partial w}{\partial y}\right).$$

By elementary calculations

$$w_z = 0 \quad \text{and} \quad w_{\overline{z}} = -\frac{1}{\overline{z}^2}.$$

Hence

$$|w'(z)| = \frac{1}{|\overline{z}|^2} \quad \text{and} \quad |J(z,w)| = \frac{1}{|\overline{z}|^4}.$$

Finally we get

$$\iint\limits_{R_\delta} |\nabla f(w(z))|\ dxdy \le \sup_{z\in R_\delta} |\overline{z}|^2 \iint\limits_{D_\delta} |\nabla f|(w)\ dudv \le \delta^2 \iint\limits_{D(0,1)} |\nabla f|(w)\ dudv.$$

In order to estimate the integral

$$\iint\limits_{A} |f\left(w(z)\right)|\ dxdy$$

we will use the change of variable formula.

We obtain

$$\begin{aligned}\iint\limits_{R_\delta} |f\left(w(z)\right)|\ dxdy &= \iint\limits_{R_\delta} |f\left(w(z)\right)||J(z,w)||J(z,w)|^{-1}\ dxdy \\ &\le \sup_{z\in R_\delta} \frac{1}{|J(z,w)|} \iint\limits_{R_\delta} |f\left(w(z)\right)||J(z,w)|\ dxdy \\ &= \sup_{z\in R_\delta} \frac{1}{|J(z,w)|} \iint\limits_{D_\delta} |f(w)|\ dudv \le \delta^4 \iint\limits_{D(0,1)} |f(w)|\ dudv.\end{aligned}$$

Using the following Poincaré–Sobolev inequality [1]

$$\iint\limits_{\Omega} |f(z)|\ dxdy \le \frac{d}{2} \iint\limits_{\Omega} |\nabla f(z)|\ dxdy$$

where Ω be a convex domain with diameter d and $f \in W_1^1(\Omega)$, finally we obtain

$$\iint\limits_{R_\delta} |f\left(w(z)\right)|\ dxdy \le \delta^4 \iint\limits_{D(0,1)} |\nabla f(w)|\ dudv.$$

Thus

$$\iint\limits_{R_\delta} |\nabla \widetilde{f}(z)|\ dxdy \le \delta^2 \iint\limits_{D(0,1)} |\nabla f(w)|\ dudv + \frac{\delta^5}{\delta - 1} \iint\limits_{D(0,1)} |\nabla f(w)|\ dudv,$$

and consequently

$$\begin{aligned}\|E(f) \mid L_1^1(D(0,\delta))\| &\le \left(1 + \delta^2 + \frac{\delta^5}{\delta - 1}\right) \iint\limits_{D(0,1)} |\nabla f(z)|\ dxdy \\ &= \left(1 + \delta^2 + \frac{\delta^5}{\delta - 1}\right) \|f \mid L_1^1(D(0,1))\|.\end{aligned}$$

Now, by the Gagliardo inequality [13, 23] we obtain

$$\left(\iint\limits_{D(0,1)} |f(z)|^2\ dxdy\right)^{\frac{1}{2}}$$

$$\leq \left(\iint\limits_{D(0,\delta)} |Ef(z)|^2\ dxdy\right)^{\frac{1}{2}} \leq \frac{1}{2\sqrt{\pi}} \iint\limits_{D(0,\delta)} |\nabla Ef(z)|\ dxdy$$

$$\leq \frac{\delta^5+\delta^3-\delta^2+\delta-1}{2\sqrt{\pi}(\delta-1)} \iint\limits_{D(0,1)} |\nabla f(z)|\ dxdy \leq C(\delta) \iint\limits_{D(0,1)} |\nabla f(z)|\ dxdy.$$

Taking the optimal $\delta = 5/4$ we have

$$\left(\iint\limits_{D(0,1)} |f(z)|^2\ dxdy\right)^{\frac{1}{2}} \leq \frac{3\sqrt{\pi^3}}{4} \iint\limits_{D(0,1} |\nabla f(z)|\ dxdy.$$

□

3. The Doubling Condition and the Hölder Inequality

Recall necessary facts about conformal capacity. A well-ordered triple $(F_0, F_1; \Omega)$ of nonempty sets, where Ω is an open set in $\mathbb{C}$, and F_0, F_1 are closed subsets of Ω, is called a condenser on the complex plane $\mathbb{C}$.

The value

$$\operatorname{cap}(E) = \operatorname{cap}(F_0, F_1; \Omega) = \inf \int\limits_{\Omega} |\nabla v|^2 dxdy,$$

where the infimum is taken over all Lipschitz nonnegative functions $v : \overline{\Omega} \to \mathbb{R}$, such that $v = 0$ on F_0, and $v = 1$ on F_1, is called conformal capacity of the condenser $E = (F_0, F_1; \Omega)$. If the set of admissible functions is empty, then $\operatorname{cap}(F_0, F_1; \Omega) = \infty$. For finite values of capacity $0 < \operatorname{cap}(F_0, F_1; \Omega) < +\infty$ there exists a unique continuous weakly differentiable function u_0 (an extremal function) such that:

$$\operatorname{cap}(F_0, F_1; \Omega) = \iint\limits_{\Omega} |\nabla u_0|^2 dxdy.$$

Quasiconformal mappings can be characterized in capacitary terms (see, for example [17]). Namely, a homeomorphism $\varphi : \Omega \to \Omega'$, $\Omega, \Omega' \subset \mathbb{C}$ is K-quasiconformal, if and only if

$$K^{-1} \operatorname{cap}(F_0, F_1; \Omega) \leq \operatorname{cap}(\varphi(F_0), \varphi(F_1); \Omega') \leq K\ \operatorname{cap}(F_0, F_1; \Omega)$$

for any condenser $E = (F_0, F_1; \Omega)$.

We will need the following estimate of the conformal capacity (see, for example [17, 29]).

Denote $D(z_0, r)$ an open disc with center z_0 of radius r and $\overline{D(z_0, r)}$ its closure.

Lemma 3.1. [17] *Let* $R > r > 0$. *Then* $\operatorname{cap}\left(\overline{D(z_0, r)}, \mathbb{C} \setminus D(z_0, R); \mathbb{C}\right) = 2\pi(\log R/r)^{-1}$.

Consider capacity estimates for Teichmüller type condensers in $\overline{\mathbb{C}}$.

Lemma 3.2. *Fix $0 < r < R$. Let F_0 and F_1 be continuums in $\mathbb{C}$ such that*

$$F_0 \cap S(0,\rho) \neq \emptyset \text{ and } F_1 \cap S(0,\rho) \neq \emptyset$$

for all $r < \rho < R$, where $S(0,\rho)$ is the circle of radius ρ. Then

$$\operatorname{cap}(F_0, F_1; D(0,R) \setminus \overline{D(0,r)}) \geq \frac{2}{\pi} \log \frac{R}{r}.$$

Denote by $R_T(t) = \overline{\mathbb{C}} \setminus \{[-1,0] \cup [t,\infty]\}$, $t > 1$, the Teichmüller ring in $\overline{\mathbb{C}}$.

Lemma 3.3. *Let $R_T(t)$ be the Teichmüller ring in $\overline{\mathbb{C}}$. Then*

$$\operatorname{cap} R_T(t) = \frac{2\pi}{\log \Phi(t)},$$

where Φ satisfies the conditions

$$t + 1 \leq \Phi(t) < 32t, \ t > 1.$$

Using this capacity estimates we obtain estimates for a constant in the inverse Hölder inequality. We start from the weak Hölder inequality.

Lemma 3.4. *Let $\varphi : \Omega \to \Omega'$ be a K-quasiconformal mapping of planar domains $\Omega, \Omega' \subset \mathbb{C}$. Then for every disc $D(z_0, r)$ such that $D(z_0, 2r) \subset \Omega$ the weak inverse Hölder inequality for the first generalized derivatives of φ*

$$\left(\frac{1}{|D(z_0,r)|} \iint\limits_{D(z_0,r)} |\varphi'(z)|^2 \, dxdy \right)^{\frac{1}{2}} \leq \frac{24\pi^2 K}{|D(z_0,2r)|} \iint\limits_{D(z_0,2r)} |\varphi'(z)| \, dxdy \tag{3.1}$$

holds.

Proof. Derivatives of K-quasiconformal mappings satisfy the following inequality ([8], pp 274)

$$\left(\frac{1}{|D(z_0,r)|} \iint\limits_{D(z_0,r)} |\varphi'(z)|^2 \, dxdy \right)^{\frac{1}{2}}$$
$$\leq \frac{4K}{r} \left(\frac{1}{|D(z_0,2r)|} \iint\limits_{D(z_0,2r)} \left| \varphi(z) - \frac{1}{|D(z_0,2r)|} \iint\limits_{D(z_0,2r)} \varphi(z) \right|^2 dxdy \right)^{\frac{1}{2}}.$$

Applying to the right side of this inequality the Poincaré-Sobolev inequality (2.1), we obtain the required inequality (3.1). □

By Lemma 3.4 and Theorem 4.2 from [8] we have the following weak Hölder inequality:

Theorem 3.5. *Let $\varphi : \Omega \to \Omega'$ be a K-quasiconformal mapping of planar domains $\Omega, \Omega' \subset \mathbb{C}$. Then for every disc $D(z_0, r)$ such that $D(z_0, 2r) \subset \Omega$ and for some*

$\sigma > 2$ the inequality

$$\left(\frac{1}{|D(z_0,r)|}\iint\limits_{D(z_0,r)} |\varphi'(z)|^\sigma \, dxdy\right)^{\frac{1}{\sigma}} \leq C_\sigma \left(\frac{1}{|D(z_0,2r)|}\iint\limits_{D(z_0,2r)} |\varphi'(z)|^2 \, dxdy\right)^{\frac{1}{2}} \tag{3.2}$$

holds, where

$$C_\sigma = \frac{10^6}{[(\sigma-1)(1-\nu)]^{1/\sigma}}, \quad \nu = 10^{4\sigma}\frac{\sigma-2}{\sigma-1}\left(24\pi^2 K\right)^\sigma < 1.$$

For estimates of the left side of the inequality (3.2) we use the doubling property of measures generated by Jacobians of quasiconformal mappings (see, for example, [21]). Recall that a Borel measure μ on a set Ω is doubling if there exist a constant $C_\mu \geq 1$ so that the inequality

$$\mu(D(z_0,2r)) \leq C_\mu \cdot \mu(D(z_0,r)) < \infty$$

hold for all discs $D(z_0,r)$ in Ω.

In the following lemma we give an estimate of the constant C_μ in the measure doubling condition.

Proposition 3.6. *Let $\varphi : \mathbb{C} \to \mathbb{C}$ be a K-quasiconformal mapping. Then for any $z_0 \in \mathbb{C}$ and any $r > 0$ we have*

$$\iint\limits_{D(z_0,2r)} |J(z,\varphi)| \, dxdy \leq \exp\left\{\frac{K\pi^2(2+\pi^2)^2}{2\log 3}\right\} \iint\limits_{D(z_0,r)} |J(z,\varphi)| \, dxdy. \tag{3.3}$$

Proof. Because the inequality (3.3) is invariant under translations and similarities we can suppose that $z_0 = 0$ and radius $r = 1$. It is sufficient to show that

$$|\varphi(D(0,2))| \leq \exp\left\{\frac{K\pi^2(2+\pi^2)^2}{2\log 3}\right\} |\varphi(D(0,1))|.$$

Since φ is a quasiconformal mapping, $\varphi(0)$ is an interior point of the open connected set $U = \varphi(D(0,1))$. Denote by $\lambda_0 = \operatorname{dist}(\varphi(0), \partial U)$ and let

$$\lambda_1 = \max_{y \in \partial \widetilde{U}} |\varphi(0) - y|, \text{ where, } \widetilde{U} = \varphi(D(0,2)).$$

Then $2\lambda_1 \geq \operatorname{diam}(\widetilde{U})$. Next let F_0 be a line segment of length λ_0 joining $\varphi(0)$ to ∂U and F_1 be a continuum in $\overline{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ joining a point in $\partial D(\varphi(0), \lambda_1) \cap \widetilde{U}$ to ∞ in $\overline{\mathbb{C}} \setminus D(\varphi(0), \lambda_1)$.

Denote by w_0 the point of intersection of F_0 and ∂U. Then $z_0 = \varphi^{-1}(w_0) = (1, \theta_0)$, where $(1, \theta_0)$ are the polar coordinates of the point z_0.

Consider the $\psi : \mathbb{R}^2 \to \mathbb{R}^2$ defined by

$$\psi(r,\theta) = (r, \theta - (r-1)\theta_0),\ 0 \leq \theta_0 \leq \pi.$$

The transformation ψ is a bi-Lipschtz mapping with the Lipschitz coefficient $L \leq \sqrt{(2+\pi^4)/2}$.

We denote by R the set $D(\frac{3}{2},\frac{3}{2}) \setminus D(\frac{3}{2},\frac{1}{2})$. Then by Lemma 3.2 using the monotonicity of the capacity and quasi-invariance under bi-Lipschtz mappings we obtain

$$\operatorname{cap}(\varphi^{-1}(F_0),\varphi^{-1}(F_1);D(0,\pi)) \geq \frac{1}{L^4}\operatorname{cap}(E_0,E_1;R) \geq \frac{8\log 3}{\pi(2+\pi^4)^2}, \tag{3.4}$$

where $E_0 = \psi(\varphi^{-1}(F_0)) \cap R$, $E_1 = \psi(\varphi^{-1}(F_1)) \cap R$.

On the other hand, taking into account the capacitary definition of the quasiconformal mapping and Lemma 3.1 we have

$$\begin{aligned}\operatorname{cap}(\varphi^{-1}(F_0),\varphi^{-1}(F_1);D(0,\pi)) &\leq K\operatorname{cap}(F_0,F_1;\mathbb{C}) \\ &\leq K\operatorname{cap}(D(\varphi(0),\lambda_0),\mathbb{C}\setminus D(\varphi(0),\lambda_1);\mathbb{C}) = \frac{2\pi K}{\log\frac{\lambda_1}{\lambda_0}}.\end{aligned} \tag{3.5}$$

Combining inequalities (3.4) and (3.5), we obtain

$$\frac{8\log 3}{\pi(2+\pi^4)^2} \leq \frac{2\pi K}{\log\frac{\lambda_1}{\lambda_0}}.$$

By elementary calculations

$$\lambda_1 \leq \exp\left\{\frac{K\pi^2(2+\pi^2)^2}{4\log 3}\right\}\lambda_0.$$

In order to obtain the required inequality necessary perform the following straightforward calculations

$$\begin{aligned}|\varphi(D(0,2))| &\leq \pi\lambda_1^2 \\ &\leq \pi\exp\left\{\frac{K\pi^2(2+\pi^2)^2}{2\log 3}\right\}\lambda_0^2 \leq \exp\left\{\frac{K\pi^2(2+\pi^2)^2}{2\log 3}\right\}|\varphi(D(0,1))|.\end{aligned}$$

Thus

$$\iint\limits_{D(0,2)} |J(z,\varphi)|\,dxdy \leq \exp\left\{\frac{K\pi^2(2+\pi^2)^2}{2\log 3}\right\}\iint\limits_{D(0,1)} |J(z,\varphi)|\,dxdy.$$

□

For any planar K-quasiconformal homeomorphism $\varphi : \Omega \to \Omega'$ the following sharp result is known: $J(z,\varphi) \in L_{p,\mathrm{loc}}(\Omega)$ for any $1 \leq p < \frac{K}{K-1}$ ([5, 16]).

Proposition 3.7. *For any conformal homeomorphism $\varphi : \mathbb{D} \to \Omega$ of the unit disc $\mathbb{D}$ onto a K-quasidisc Ω derivatives $\varphi' \in L_p(\mathbb{D})$ for any $1 \leq p < \frac{2K^2}{K^2-1} \subset \mathrm{HI}(\Omega)$.*

Proof. Any conformal homeomorphism $\varphi : \mathbb{D} \to \Omega$ can be extended to a K^2 quasiconformal homeomorphism $\widetilde{\varphi}$ of the whole plane to the whole plane by reflection. Hence $\widetilde{\varphi}'$ belongs to the class $L_{p,\mathrm{loc}}(\mathbb{C})$ for any $1 \leq p < \frac{2K^2}{K^2-1}$ ([5, 16]). Therefore φ' belongs to the class $L_p(\mathbb{D})$. □

On the base of the weak Hölder inequality and the doubling condition we obtain integral estimates of complex derivatives of conformal mapping $\varphi : \mathbb{D} \to \Omega$ in the unit disc onto a K-quasidisc Ω:

Theorem B. *Let $\Omega \subset \mathbb{R}^2$ be a K-quasidisc and $\varphi : \mathbb{D} \to \Omega$ be a conformal mapping. Suppose that $2 < \alpha < \frac{2K^2}{K^2-1}$. Then*

$$\left(\iint\limits_{D(0,1)} |J(z,\varphi)|^{\frac{\alpha}{2}} \, dxdy \right)^{\frac{2}{\alpha}} \leq \frac{C_\alpha^2 K^2 \pi^{\frac{2}{\alpha}-1}}{4} \exp\left\{ \frac{K^2\pi^2(2+\pi^2)^2}{2\log 3} \right\} \cdot |\Omega|. \tag{3.6}$$

where

$$C_\alpha = \frac{10^6}{[(\alpha-1)(1-\nu)]^{1/\alpha}}, \quad \nu = 10^{4\alpha}\frac{\alpha-2}{\alpha-1}(24\pi^2K^2)^\alpha < 1.$$

Proof. Since $\varphi : \mathbb{D} \to \Omega$ is a conformal mapping, then by the inequality (3.2) using the equality $J(z,\varphi) = |\varphi'(z)|^2$ we have

$$\left(\iint\limits_{D(0,1)} |J(z,\varphi)|^{\frac{\alpha}{2}} \, dxdy \right)^{\frac{2}{\alpha}} = |D(0,1)|^{\frac{2}{\alpha}} \left(\frac{1}{|D(0,1)|} \iint\limits_{D(0,1)} |\varphi'(z)|^\alpha \, dxdy \right)^{\frac{1}{\alpha}\cdot 2}$$

$$\leq C_\alpha^2 |D(0,1)|^{\frac{2}{\alpha}} \left(\frac{1}{|D(0,2)|} \iint\limits_{D(0,2)} |\varphi'(z)|^2 \, dxdy \right) = \frac{C_\alpha^2 \pi^{\frac{2}{\alpha}-1}}{4} \iint\limits_{D(0,2)} |\varphi(z)|^2 \, dxdy.$$

In the disc $D(0,2)$ an extension of the conformal mapping φ is a K^2-quasiconformal homeomorphism. Hence taking into account the inequality

$$|\varphi'(z)|^2 \leq K^2 |J(z,\varphi)| \ \text{ for almost all } z \in D(0,2)$$

by the inequality (3.3) we get

$$\frac{C_\alpha^2 \pi^{\frac{2}{\alpha}-1}}{4} \iint\limits_{D(0,2)} |\varphi'(z)|^2 \, dxdy \leq \frac{C_\alpha^2 K^2 \pi^{\frac{2}{\alpha}-1}}{4} \iint\limits_{D(0,2)} |J(z,\varphi)| \, dxdy$$

$$\leq \frac{C_\alpha^2 K^2 \pi^{\frac{2}{\alpha}-1}}{4} \exp\left\{ \frac{K^2\pi^2(2+\pi^2)^2}{2\log 3} \right\} \iint\limits_{D(0,1)} |J(z,\varphi)| \, dxdy.$$

Thus, considering that

$$\iint\limits_{D(0,1)} |J(z,\varphi)| \, dxdy = |\Omega|$$

we have

$$\left(\iint\limits_{D(0,1)} |J(z,\varphi)|^{\frac{\alpha}{2}} \, dxdy \right)^{\frac{2}{\alpha}} \leq \frac{C_\alpha^2 K^2 \pi^{\frac{2}{\alpha}-1}}{4} \exp\left\{ \frac{K^2\pi^2(2+\pi^2)^2}{2\log 3} \right\} \cdot |\Omega|.$$

□

4. Estimates of the hyperbolic α-dilatation and the first non-trivial eigenvalue of the Neumann–Laplace operator

In the work [19] was obtained the following estimates of the first non-trivial eigenvalue of the Neumann-Laplace operator in quasidiscs:

Proposition 4.1. *Suppose a conformal homeomorphism $\varphi : \mathbb{D} \to \Omega$ maps the unit disc $\mathbb{D}$ onto a K-quasidisc Ω. Then*

$$1/\mu_1(\Omega) \leq \frac{4}{\sqrt[\alpha]{\pi^2}} \left(\frac{2\alpha-2}{\alpha-2}\right)^{\frac{2\alpha-2}{\alpha}} Q(\alpha,\Omega)^{2/\alpha}$$

for any $2 < \alpha < \frac{2K^2}{K^2-1}$.

By Proposition 4.1 and Theorem B we obtain the lower estimates of the first non-trivial eigenvalue of the Neumann–Laplace operator in K-quasidisc in terms of quasiconformal geometry of domains:

Theorem C. *Let $\Omega \subset \mathbb{C}$ be a K-quasidisc. Then the spectrum of Neumann-Laplace operator in Ω is discrete, can be written in the form of a non-decreasing sequence*

$$0 = \mu_0(\Omega) < \mu_1(\Omega) \leq \mu_2(\Omega) \leq ... \leq \mu_n(\Omega) \leq ...,$$

and

$$\frac{1}{\mu_1(\Omega)} \leq \frac{K^2 C_\alpha^2}{\pi} \left(\frac{2\alpha-2}{\alpha-2}\right)^{\frac{2\alpha-2}{\alpha}} \exp\left\{\frac{K^2\pi^2(2+\pi^2)^2}{2\log 3}\right\} \cdot \big|\Omega\big|, \tag{4.1}$$

for $2 < \alpha < \frac{2K^2}{K^2-1}$, where

$$C_\alpha = \frac{10^6}{[(\alpha-1)(1-\nu)]^{1/\alpha}}, \quad \nu = 10^{4\alpha}\frac{\alpha-2}{\alpha-1}(24\pi^2K^2)^\alpha < 1.$$

Proof. By Proposition 4.1 for any $2 < \alpha < \frac{2K^2}{K^2-1}$ we have

$$1/\mu_1(\Omega) \leq \frac{4}{\sqrt[\alpha]{\pi^2}} \left(\frac{2\alpha-2}{\alpha-2}\right)^{\frac{2\alpha-2}{\alpha}} Q(\alpha,\Omega)^{2/\alpha}. \tag{4.2}$$

By Theorem B

$$Q\,(\alpha,\Omega)^{\frac{2}{\alpha}} := \left(\iint\limits_{D(0,1)} |J(z,\varphi)|^{\frac{\alpha}{2}}\, dxdy\right)^{\frac{2}{\alpha}} \leq \frac{C_\alpha^2 K^2 \pi^{\frac{2}{\alpha}-1}}{4} \exp\left\{\frac{K^2\pi^2(2+\pi^2)^2}{2\log 3}\right\} \cdot |\Omega|. \tag{4.3}$$

Combining inequalities (4.2) and (4.3) and performing a straightforward calculations, we obtain the required inequality. □

As an application of Theorem C, we obtain the lower estimates of the first non-trivial eigenvalue on the Neumann eigenvalue problem for the Laplace operator in the star-shaped and spiral-shaped domains.

Star-shaped domains. We say that a domain Ω^* is β-star-shaped (with respect to $z_0 = 0$) if the function $\varphi(z)$, $\varphi(0) = 0$, conformally maps a unit disc $\mathbb{D}$ onto Ω^* and the condition satisfies [12]:

$$\left|\arg \frac{z\varphi'(z)}{\varphi(z)}\right| \le \beta\pi/2, \quad 0 \le \beta < 1, \quad |z| < 1.$$

In [12] proved the following: the boundary of the β-star-shaped domain Ω^* is a K-quasicircle with $K = \cot^2(1-\beta)\pi/4$.

Then by Theorem C we have

$$\frac{1}{\mu_1(\Omega^*)} \le \frac{C_\alpha^2 \cot^4(1-\beta)\frac{\pi}{4}}{\pi}\left(\frac{2\alpha-2}{\alpha-2}\right)^{\frac{2\alpha-2}{\alpha}} \exp\left\{\frac{\pi^2(2+\pi^2)^2\cot^4(1-\beta)\frac{\pi}{4}}{2\log 3}\right\}\cdot\left|\Omega^*\right|,$$

where

$$C_\alpha = \frac{10^6}{[(\alpha-1)(1-\nu)]^{1/\alpha}}, \quad \nu = 10^{4\alpha}(24\pi^2\cot^4(1-\beta)\pi/4)^\alpha\frac{\alpha-2}{\alpha-1} < 1.$$

Spiral-shaped domains. We say that a domain Ω_s is β-spiral-shaped (with respect to $z_0 = 0$) if the function $\varphi(z)$, $\varphi(0) = 0$, conformally maps a unit disc $\mathbb{D}$ onto Ω_s and the condition satisfies [27]:

$$\left|\arg e^{i\delta}\frac{z\varphi'(z)}{\varphi(z)}\right| \le \beta\pi/2, \quad 0 \le \beta < 1, \quad |\delta| < \beta\pi/2, \quad |z| < 1.$$

In [27] proved the following: the boundary of the β-spiral-shaped domain Ω_s is a K-quasicircle with $K = \cot^2(1-\beta)\pi/4$.

Then by Theorem C we have

$$\frac{1}{\mu_1(\Omega_s)} \le \frac{C_\alpha^2 \cot^4(1-\beta)\frac{\pi}{4}}{\pi}\left(\frac{2\alpha-2}{\alpha-2}\right)^{\frac{2\alpha-2}{\alpha}} \exp\left\{\frac{\pi^2(2+\pi^2)^2\cot^4(1-\beta)\frac{\pi}{4}}{2\log 3}\right\}\cdot\left|\Omega_s\right|,$$

where

$$C_\alpha = \frac{10^6}{[(\alpha-1)(1-\nu)]^{1/\alpha}}, \quad \nu = 10^{4\alpha}(24\pi^2\cot^4(1-\beta)\pi/4)^\alpha\frac{\alpha-2}{\alpha-1} < 1.$$

5. Quasiconformal reflection

Let Γ be a Jordan curve on the Riemann sphere, and denote its complementary components by Ω, Ω^*. Suppose that there exists a sense-reversing quasiconformal mapping ψ of the sphere onto itself which maps Ω on Ω^* and keeps every point on Γ fixed. Such mappings are called quasiconformal reflections.

Denote by H the upper half-plane and by H^* the lower half-plane of the complex plane $\mathbb{C}$. Consider a conformal mapping φ of H on Ω and a conformal mapping φ_* of H^* to Ω^*. Then a mapping $\varphi_*^{-1}\circ\psi\circ\varphi$ is a quasiconformal mapping of H onto H^* which induces a monotone mapping $h = \varphi_*^{-1}\circ\varphi$ of the real axis on itself [2].

In [2] L. Ahlfors and A. Beurling derived a necessary and sufficient condition for a boundary mapping h to be restriction of a quasiconformal mapping of H on itself (or on its reflection H^*). Without loss of generality it may be assumed that

$h(\infty) = \infty$. Then h admits a quasiconformal extension if and only if it satisfies a M-condition, namely an inequality

$$\frac{1}{M} \le \frac{h(x+t) - h(x)}{h(x) - h(x-t)} \le M \tag{5.1}$$

which is to be fulfilled for all real x, t, $t > 0$, and with a constant $M \ne 0, \infty$. More precisely, if h has a K-quasiconformal extension, then (5.1) holds with a $M = M(K) < e^{\pi K}/16$, and if (5.1) holds, then h has a K-quasiconformal extension such that $K = K(M) < M^2$.

In [3] L. Ahlfors proved that a Jordan curve Γ admits a K-quasiconformal reflection if and only if Γ satisfies the Ahlfors's 3-point condition. In the following theorem, using the Ahlfors scheme, we calculates the upper estimates for the coefficient of quasiconformality K for quasicircles given by the Ahlfors's 3-point condition.

Theorem 5.1. *Let a Jordan curve Γ satisfies the Ahlfors's 3-point condition: there exists a constant C such that*

$$|\zeta_3 - \zeta_1| \le C|\zeta_2 - \zeta_1| \tag{5.2}$$

for any three point on Γ where ζ_3 is between ζ_1 and ζ_2. Then Γ to admit a K-quasiconformal reflection where K depends only on C and

$$K < \frac{1}{2^{10}} \exp\left\{\left(1 + e^{2\pi} C^5\right)^2\right\}.$$

Proof. We will use the notations

$$\alpha_1 = \operatorname{arc} \zeta_1\zeta_3, \quad \alpha_2 = \operatorname{arc} \zeta_3\zeta_2, \quad \beta_1 = \operatorname{arc} \zeta_2\infty, \quad \beta_2 = \operatorname{arc} \zeta_1\infty.$$

Thus

$$\operatorname{cap}(\alpha_1, \beta_1; \Omega) \cdot \operatorname{cap}(\alpha_2, \beta_2; \Omega) = 1 \text{ and } \operatorname{cap}(\alpha_1, \beta_1; \Omega^*) \cdot \operatorname{cap}(\alpha_2, \beta_2; \Omega^*) = 1.$$

Through the conformal mapping of Ω, let ζ_1, ζ_3, ζ_2 correspond to $x - t$, x, $x + t$. This means that

$$\operatorname{cap}(\alpha_1, \beta_1; \Omega) = \operatorname{cap}(\alpha_2, \beta_2; \Omega) = 1.$$

Through the conformal mapping of Ω^*, ζ_1, ζ_3, ζ_2 correspond to $h(x - t)$, $h(x)$, $h(x + t)$.

In [3] L. Ahlfors prove that

$$\operatorname{cap}(\alpha_1, \beta_1; \Omega^*) \le \pi \left(1 + e^{2\pi} C^5\right)^2. \tag{5.3}$$

Now we obtain the lower estimate for the capacity $\operatorname{cap}(\alpha_1, \beta_1; \Omega^*)$ using the capacity estimates for the Teichmüller condenser: denote by

$$y = (h(x+t) - h(x))/(h(x) - h(x-t)),$$

then by Lemma 3.3

$$\operatorname{cap} R_T(y) = \frac{2\pi}{\log \Phi(y)} > \frac{2\pi}{\log 32y}. \tag{5.4}$$

Combining inequalities (5.3) and (5.4), we obtain

$$\frac{2\pi}{\log 32y} < \operatorname{cap} R_T(y) = \operatorname{cap}(\alpha_1, \beta_1; \Omega^*) \le \pi\left(1 + e^{2\pi} C^5\right)^2$$

or

$$\frac{h(x+t) - h(x)}{h(x) - h(x-t)} \le \frac{1}{32} \exp\left\{\frac{\left(1 + e^{2\pi} C^5\right)^2}{2}\right\}.$$

Hence h satisfies an M-condition, i.e.

$$\frac{1}{M(C)} \leq \frac{h(x+t)-h(x)}{h(x)-h(x-t)} \leq M(C)$$

where

$$M(C) = \frac{1}{32}\exp\left\{\frac{\left(1+e^{2\pi}C^5\right)^2}{2}\right\}.$$

So, [2] there exist a K-quasiconformal reflection such that $K < M^2(C)$.

Finally we get

$$K < \left(\frac{1}{32}\exp\left\{\frac{\left(1+e^{2\pi}C^5\right)^2}{2}\right\}\right)^2 = \frac{1}{2^{10}}\exp\left\{\left(1+e^{2\pi}C^5\right)^2\right\}. \tag{5.5}$$

□

Combining Theorem C and Theorem 5.1 we obtain

Corollary 5.2. *Let a domain $\Omega \subset \mathbb{R}^2$ is bounded by a Jordan curve Γ satisfies the Ahlfors's 3-point condition. Then*

$$\frac{1}{\mu_1(\Omega)} \leq \frac{C_\alpha^2 e^{2\left(1+e^{2\pi}C^5\right)}}{2^{20}\pi}\left(\frac{2\alpha-2}{\alpha-2}\right)^{\frac{2\alpha-2}{\alpha}}\exp\left\{\frac{\pi^2(2+\pi^2)^2 e^{2\left(1+e^{2\pi}C^5\right)}}{2^{21}\log 3}\right\}\cdot\left|\Omega\right|,$$

for $2 < \alpha < \frac{2K^2}{K^2-1}$, where

$$C_\alpha = \frac{10^6}{[(\alpha-1)(1-\nu)]^{1/\alpha}}, \quad \nu = 10^{4\alpha}\left(\frac{3\pi^2}{2^{17}}e^{2\left(1+e^{2\pi}C^5\right)}\right)^\alpha \frac{\alpha-2}{\alpha-1} < 1.$$

6. Examples in fractal type domains

Rohde snowflake. In [26] S. Rohde constructed a collection S of snowflake type planar curves with the intriguing property that each planar quasicircle is bi-Lipschitz equivalent to some curve in S.

Rohde's catalog is

$$S := \bigcup_{1/4\leq p<1/2} S_p$$

where p is a snowflake parameter. Each curve in S_p is built in a manner reminiscent of the construction of the von Koch snowflake. Thus, each $S \in S_p$ is the limit of a sequence S^n of polygons where S^{n+1} is obtained from S^n by using the replacement rule illustrated in Figure 6.1: for each of the 4^n edges E of S^n we have two choices, either we replace E with the four line segments obtained by dividing E into four arcs of equal diameter, or we replace E by a similarity copy of the polygonal arc A_p pictured at the top right of Figure 6.1. In both cases E is replaced by four new segments, each of these with diameter $(1/4)\mathrm{diam}(E)$ in the first case or with diameter $p\,\mathrm{diam}(E)$ in the second case. The second type of replacement is done so that the "tip" of the replacement arc points into the exterior of S^n. This iterative process starts with S^1 being the unit square, and the snowflake parameter, thus the polygon arc A_p, is fixed throughout the construction.

The sequence S^n of polygons converges, in the Hausdorff metric, to a planar quasicircle S that we call a *Rohde snowflake* constructed with snowflake parameter p. Then S_p is the collection of all Rohde snowflakes that can be constructed with snowflake parameter p.



FIGURE 6.1. Construction of a Rohde-snowflake.

In [22] established that each Rohde snowflake S in S_p is C-bounded turning with

$$C = C(p) = \frac{16}{1-2p}, \quad 1/4 \le p < 1/2.$$

A planar curve Γ satisfies the C-bounded turning, $C \ge 1$, if for each pair of points x, y, on Γ, the smaller diameter subarc $\Gamma[x,y]$ of Γ that joins x, y satisfies

$$\operatorname{diam}(\Gamma[x,y]) \le C|x-y|. \tag{6.1}$$

The C-bounded turning condition (6.1) is equivalent the Ahlfors's 3-point condition (5.2) with the same constant C [17].

The following theorem gives the lower estimates of the first non-trivial eigenvalue of the Neumann–Laplace operator in domains type a Rohde snowflakes:

Theorem A. *Let $S_p \subset \mathbb{R}^2$, $1/4 \le p < 1/2$, be the Rohde snowflake. Then the spectrum of Neumann-Laplace operator in S_p is discrete, can be written in the form of a non-decreasing sequence*

$$0 = \mu_0(S_p) < \mu_1(S_p) \le \mu_2(S_p) \le \ldots \le \mu_n(S_p) \le \ldots,$$

and

$$\frac{1}{\mu_1(S_p)} \le \frac{C_\alpha^2 e^{4\left(1+e^{2\pi}(16/(1-2p))^5\right)}}{2^{40}\pi} \left(\frac{2\alpha-2}{\alpha-2}\right)^{\frac{2\alpha-2}{\alpha}} \times \exp\left\{\frac{\pi^2(2+\pi^2)^2 e^{4\left(1+e^{2\pi}(16/(1-2p))^5\right)}}{2^{41}\log 3}\right\} |S_p|,$$

for $2 < \alpha < \frac{2K^2}{K^2-1}$, where

$$C_\alpha = \frac{10^6}{[(\alpha-1)(1-\nu)]^{1/\alpha}}, \quad \nu = 10^{4\alpha}\left(\frac{3\pi^2}{2^{37}} e^{4\left(1+e^{2\pi}C^5\right)}\right)^\alpha \frac{\alpha-2}{\alpha-1} < 1.$$

The proof Theorem A immediately follows from the Corollary 5.2, given that any L-bi-Lipschitz planar homeomorphism is K-quasiconformal with $K = L^2$.

Appendix

Firstly we discuss a new notion of the hyperbolic α-dilatation $Q(\alpha,\Omega)$ and its convergence hyperbolic interval $\mathrm{HI}(\Omega)$ in connection with known results. Of course, quotients $Q(\alpha,\Omega)$ and $\mathrm{HI}(\Omega)$ can be defined also in non bounded domains.

Recall definitions:

$$Q(\alpha,\Omega):=\iint\limits_{\mathbb{D}}|\varphi'(z)|^{\alpha}\,dxdy=\iint\limits_{\Omega}\left|\left(\varphi^{-1}\right)'(w)\right|^{2-\alpha}dudv$$

where $\varphi:\mathbb{D}\to\Omega$ is a Riemann conformal homeomorphism;

$$\mathrm{HI}(\Omega):=\{\alpha\in\mathbb{R}:Q(\alpha,\Omega)<\infty\}.$$

Let us remark that by the definition

$$Q(2,\Omega)=|\Omega|,$$

i. e is finite for any domain of finite measure. We don't know any simple interpretation of $Q(\alpha,\Omega)$ for a number α other than 2.

In these new terms of **the Inverse Brennan's conjecture** [9] states: Let $\Omega\subset\mathbb{C}$ be a simply connected planar domain with nonempty boundary, and $\varphi:\mathbb{D}\to\Omega$ be a conformal homeomorphism. Then

$$\mathrm{HI}(\Omega)=(-2,2/3).$$

The upper bound $\alpha=2/3$ is proved, the lower bound is only conjectured.

By [18] for bounded domains it is proved that

$$\mathrm{HI}(\Omega)=(-2,2].$$

The upper bound $\alpha=2$ is proved, the lower bound is only conjectured.

Note that for smooth bounded domains [9]

$$\mathrm{HI}(\Omega)=(-\infty,\infty).$$

In [19] we demonstrated that for any K-quasidisc $1\le\alpha<\frac{2K^2}{K^2-1}\subset\mathrm{HI}(\Omega)$. It can be reformulated in terms of the Ahlfors condition.

Conjecture. Interval $\mathrm{HI}(\Omega)$ is defined by the hyperbolic geometry of Ω.

References

[1] G. Acosta, R. G. Durán, An optimal Poincaré inequality in L^1 for convex domains, Proc. Amer. Math. Soc. 132 (2004), no. 1, 195–202.

[2] L. Ahlfors, A. Beurling, The boundary correspondence under quasiconformal mapping, Acta Math., 56 (1956), 125–142.

[3] L. Ahlfors, Quasiconformal reflections, Acta Math., 109 (1963), 291–301.

[4] L. Ahlfors, Lectures on quasiconformal mappings, D. Van Nostrand Co., Inc., Toronto, Ont.-New York-London, 1966.

[5] K. Astala, Area distortion of quasiconformal mappings, Acta Math. 173 (1994), 37–60.

[6] A. F. Beardon, D. Minda, D. The hyperbolic metric and geometric function theory. Quasiconformal mappings and their applications, Narosa, New Delhi, (2007), 9–56.

[7] L. Bieberbach, Über die Koeffizienten derjenigen Polenzreihen, welche eine schlichte Abbildung des Einheitskreises vermitteln, S.-B. Preuss. Akad. Wiss. (1916) 940–955.

[8] B. Bojarski, T. Iwaniec, Analytical foundation of the theory of quasiconformal mappings in $\mathbb{R}^n$, Ann. Acad. Sci. Fenn. Ser. A. I. Math. 8 (1983), 257–324.

[9] J. Brennan, The integrability of the derivative in conformal mapping. J. London Math. Soc. (2) 18 (1978), no. 2, 261–272.

[10] V. I. Burenkov, V. Gol'dshtein, A. Ukhlov, Conformal spectral stability for the Dirichlet–Laplace operator. Math. Nachr. 288, (2015), 1822–1833.
[11] L. Esposito, C. Nitsch, C. Trombetti, Best constants in Poincaré inequalities for convex domains, J. Convex Anal., 20 (2013), 253–264.
[12] M. Fait, Y. Krzyz, Y. Zygmunt, Explicit quasiconformal extension for some classes of univalent functions. Comment, math, helv., 51(2), (1976), 279–285.
[13] H. Federer, W. H. Fleming, Normal and integral currents, Ann. Math., 72, (1960), 458–520.
[14] V. Ferone, C. Nitsch and C. Trombetti, A remark on optimal weighted Poincaré inequalities for convex domains, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. 23, (2012), 467–475.
[15] D. Gilbarg, N. S. Trudinger, Elliptic Partial Differential Equations of Second Order, Springer-Verlag Berlin-Heidelberg-New York (1977)
[16] V. M. Gol'dshtein, The degree of summability of generalized derivatives of quasiconformal homeomorphisms. Sibirsk. Matem. Zh. 22 (1981), no. 6, 22–40 (in Russian). (English transl.: Siberian Math. J. 22 (1981), no. 6, 821-836).
[17] V.M.Gol'dshtein, Yu.G.Reshetnyak. Quasiconformal mappings and Sobolev spaces. Kluwier Academic Publisher, Dordrecht/Boston/London, 1990.
[18] V. Gol'dshtein, A. Ukhlov, Sobolev homeomorphisms and Brennan's conjecture, Comput. Methods Funct. Theory, 14 (2014), 247–256.
[19] V. Gol'dshtein, A. Ukhlov, On the first Eigenvalues of Free Vibrating Membranes in Conformal Regular Domains. Arch. Rational Mech. Anal., 221 (2016), no. 2, 893–915.
[20] H. Hedenmalm, S. Shimorin, Weighted Bergman spaces and the integral means spectrum of conformal mappings. Duke Math. J. 127 (2005), no. 2, 341–393.
[21] J. Heinonen, P. Koskela, Weighted Sobolev and Poincaré inequalities and quasiregular mappings of polynomial type, Math. Scand., 77, (1995), 251-271.
[22] D. A. Herron, D. Meyer, Quasicircles and bounded turning circles modulo bi-Lipschitz maps, Rev. Mat. Iberoamericana, 28(3), (2012), 603-630.
[23] V. Maz'ya, Sobolev spaces: with applications to elliptic partial differential equations, Springer, Berlin/Heidelberg, 2010.
[24] L. E. Payne, H. F. Weinberger, An optimal Poincaré inequality for convex domains, Arch. Rat. Mech. Anal., 5, 286-292 (1960).
[25] G Pólya, On the eigenvalues of vibrating membranes, Proc. London Math. Soc. 11, 419–433 (1961)
[26] S. Rohde, Quasicircles modulo bi-Lipschitz maps, Rev. Mat. Iberoamericana 17 (2001), 643–659.
[27] M. A. Sevodin, Univalence conditions in spiral domains, Tr. Semin. Kraev. Zad., 23, (1986), 193–200 (in Russian).
[28] D. Valtorta, Sharp estimate on the first eigenvalue of the p-Laplacian, Nonlin. Analysis 75 (2012), 4974–4994.
[29] M. Vuorinen, Conformal geometry and quasiregular mappings, Lecture Notes in Math. Vol. 1319, Springer-Verlag, Berlin-Heidelberg-New York, 2006.

Department of Mathematics, Ben-Gurion University of the Negev, P.O.Box 653, Beer Sheva, 8410501, Israel
E-mail address: vladimir@math.bgu.ac.il

Department of Higher Mathematics and Mathematical Physics, Tomsk Polytechnic University, 634050 Tomsk, Lenin Ave. 30, Russia; Department of General Mathematics, Tomsk State University, 634050 Tomsk, Lenin Ave. 36, Russia
Current address: Department of Mathematics, Ben-Gurion University of the Negev, P.O.Box 653, Beer Sheva, 8410501, Israel
E-mail address: vpchelintsev@vtomske.ru

Department of Mathematics, Ben-Gurion University of the Negev, P.O.Box 653, Beer Sheva, 8410501, Israel
E-mail address: ukhlov@math.bgu.ac.il